



Developing a Restful Environment Action Manual:

DREAM Toolkit

Implementation Guide

This guide to the Developing a Restful Environment Action Manual (DREAM) Toolkit will help you get started with improving sleep quality for residents living with dementia.

Who can use the DREAM Toolkit?

LEADERSHIP AND SLEEP CHAMPION		ALL STAFF		INTERDISCIPLINARY TEAM	
Implementation Guide	Sleep Environment Improvement Tool	Sleep Matters Video	Sleep Handbook	DREAM Toolkit Pocket Guide	Resident Preferences Tool

Welcome to the DREAM Toolkit Implementation Guide


What is in the DREAM Toolkit?

The Developing a Restful Environment Action Manual (DREAM) Toolkit offers educational information and tools for nursing home teams to use together to improve resident sleep and quality of life.

Why does sleep matter?

Sleep helps all of us stay healthy, regardless of age. **A resident living with dementia is likely to experience worse physical, mental, and psychosocial well-being if they do not sleep well at night, creating a greater risk of adverse events, including falls.** Your team can take action today to help residents obtain higher quality, uninterrupted sleep. Develop a healthy sleep environment for **residents living with dementia** to promote safety, encourage a quality balance between wake and sleep, and enhance physical and mental health. Use the steps in this guide to get started.

Who can use the DREAM Toolkit?



How can I get started?

This guide will walk you through the following steps to get started with using the DREAM Toolkit:

1. Choose a champion.
2. Identify key areas to improve sleep.
3. Educate and empower staff.
4. Communicate the plan to staff.
5. Reach out to residents and their loved ones.
6. Schedule regular check-ins and track your progress.

A cartoon illustration of a woman with dark skin, brown hair in a bun, and glasses. She is wearing a light blue long-sleeved shirt and dark blue pants. She is giving a thumbs up with her right hand and has a friendly smile.

When Choosing a Sleep Champion:

-

Use the **Sleep Environment Improvement Tool** to decide where to focus improvement efforts and how to get started. This tool includes a Sleep Environment Scan and also offers practical approaches to improve sleep, organized by key area (e.g., light, noise, workflow).

[illegible]

Everyone Can Make a Difference:

- DREAM Toolkit Implementation Guide 2

Step 4: Communicate the Plan to Staff

Once you and your team decide on an approach, develop a plan to communicate important information to staff. Organize your communication strategy by thinking about who needs to know what. Then decide when and how to communicate the information. Consider these questions as you develop your communication plan:

- **Which departments will be involved** in the success of the approach(es)?
 - Examples: dietary, housekeeping, nursing, activities, social work, rehabilitation
- **How do they prefer to receive new information?**
 - Examples: email, in-person meeting, small group huddles, one-on-one discussion
- **Which messages are important to share?** Can any of the other components of the DREAM Toolkit support your efforts to share this information?
 - Example: the Sleep Matters video
- **When do you need to communicate this information?** Communicate before you start the approach. Then decide when it is important to check-in and review progress or concerns from your staff.
 - Examples: care planning meetings, weekly or monthly staff meetings, huddles
- **What resources can you provide for staff** to support them in rolling out the approach(es)?
 - Examples: materials on the tables in the breakroom, a binder of resources



Plan to regularly reinforce with staff the benefits of improving sleep for residents living with dementia. This will be critical for gaining staff buy-in and genuine engagement. Develop a partnership with your staff so that you can work together to troubleshoot unexpected challenges that come up as you create a better sleep environment for residents.

Step 5: Reach Out to Residents and Their Loved Ones

Use this opportunity to tell residents and their loved ones about the exciting plans to improve sleep and quality of life. You will also want to include them as partners to improve sleep and ask about resident sleep preferences.

Share and Request Information:

- Print and distribute “**We are Working to Improve Sleep for Residents**” on the next page, or create your own materials.
- Use the **Resident Preferences Tool** to collect information.
- Request ongoing input and feedback.




Each resident is the number one expert on their own preferences. Ask and engage them first. However, it may be difficult for the resident living with dementia to communicate their preferences and needs. Engage the resident's support system, including loved ones and care providers, to better understand the resident's preferences and needs. Loved ones include representatives of residents living with dementia, such as family, caregivers, friends, and court-appointed guardians. Loved ones may or may not have legal decision-making authority.

We are Working to Improve Sleep for Residents

Why focus on sleep?

Getting older, especially when living with dementia, can make it harder to get to sleep and stay asleep at night. Improving sleep can improve health and lead to better quality of life.



When residents sleep well it:

- Prevents daytime sleepiness so the resident can stay active and engaged
- Lowers risk of falls
- Improves mood



When residents have trouble sleeping it:

- Speeds up memory loss
- Reduces self-care skills
- Increases risk of infection
- Makes residents feel more confused

Medicine is not always the answer.

Your care team works together to determine when medications are appropriate. Psychotropic medications can actually cause sleep issues like insomnia and drowsiness. With long-term use, these medications can also lead to disorientation, memory loss, falls, and increased risk of worsening heart disease. Non-pharmacological approaches for sleep, such as reducing bedroom light and noise, present fewer risks than psychotropic medications.

What are we doing to improve sleep?

We are taking action to help each resident sleep better by focusing on personal sleep and wake preferences, as well as improving the sleep environment throughout the nursing home.



Help us understand what makes it easier for the resident you know to sleep well. Care planning meetings are a great time to share the following information:

- What did their bedtime routine look like at home?
- What comfort items help them sleep?
- What activities might they enjoy during the day?

Step 6: Schedule Regular Check-ins and Track Your Progress

You can measure sleep quality for residents living with dementia directly with fitness trackers (such as bracelet trackers or phone sleep apps). Data that you already use or have access to can also help you **track the progress of sleep improvement by focusing on desired outcomes**. Promoting sleep for residents living with dementia is an ongoing process. Measurable improvement will not happen immediately, but you can look for improvements over time.

Use the steps and table below to get started:

1. Identify your goals and outcomes.
2. Once you know the outcomes you want, find data sources you can use. See the table below for ideas.
3. Figure out your starting point, or baseline. Knowing where you started will make it easier to see improvement over time.
4. Continue to collect and compare data. Share the data with staff during **regular check-ins**.
5. Look at the data to learn what works and does not work in your nursing home. Use this information to adjust future plans to improve sleep quality for residents living with dementia.

Improved Sleep Outcomes	Data Sources
Increase in sleep time for residents	<ul style="list-style-type: none">• Fitness trackers worn by residents• Staff observation and daily care notes
Decrease in falls and falls with major injury	<ul style="list-style-type: none">• Internal tracking• Discussion with nurses and nurse aides about restorative care
Decrease in illness and infections	<ul style="list-style-type: none">• Internal tracking• Updates from infection preventionist, wound care specialists, and respiratory therapists
Fewer challenges in communicating unmet needs or discomfort	<ul style="list-style-type: none">• Internal tracking• Staff observation and daily care notes
Increase in resident and loved one satisfaction	<ul style="list-style-type: none">• Resident and family satisfaction surveys• Staff observation and daily care notes
Increase in staff satisfaction	<ul style="list-style-type: none">• Employee satisfaction surveys• Face-to-face feedback

Get Started with the DREAM Toolkit!

Use this Implementation Guide and other sections of the DREAM Toolkit to plan and make changes throughout your nursing home and for individual residents.



References

- Besedovsky, L., Lange, T., & Haack, M. (2019). [The Sleep-Immune Crosstalk in Health and Disease](#). *Physiological reviews*, 99(3), 1325-1380.
- Chattu, V. K., Chattu, S. K., Burman, D., Spence, D. W., & Pandi-Perumal, S. R. (2019). [The Interlinked Rising Epidemic of Insufficient Sleep and Diabetes Mellitus](#). *Healthcare (Basel, Switzerland)*, 7(1).
- Chen, J. C., Espeland, M.A., Brunner, R.L., Lovato, L.C. Wallace, R.B., Leng, X., et al. (2016). [Sleep Duration, Cognitive Decline, and Dementia Risk in Older Women](#). *Alzheimer's & Dementia*, 12(1), 21-33.
- Doghramji, K., & Jangro, W. C. (2016). [Adverse Effects of Psychotropic Medications on Sleep](#). *Psychiatric Clinics*, 39(3), 487-502.
- Furtado, F., Bruno da Silva, B. G., Abranches, I. L. L., Abrantes, A. F., & Forner-Cordero, A. (2016). [Chronic Low Quality Sleep Impairs Postural Control in Healthy Adults](#). *PLOS One*, 11(10), e0163310.
- Hennawy, M., Sabovich, S., Liu, C.S., Herrmann, N., Lanctôt K.L. (2019). [Sleep and Attention in Alzheimer's Disease](#). *Yale Journal of Biological Medicine*, 25;92(1):53-61.
- Holth, J., Patel, T., & Holtzman, D. M. (2017). [Sleep in Alzheimer's Disease - Beyond Amyloid](#). *Neurobiology Of Sleep And Circadian Rhythms*, 2, 4–14.
- Hshieh, T. T., Yue, J., Oh, E., Puelle, M., Dowal, S., Travison, T., & Inouye, S. K. (2015). [Effectiveness of Multicomponent Nonpharmacological Delirium Interventions: a Meta-Analysis](#). *JAMA Internal Medicine*, 175(4), 512–520.
- Leng, Y., Cappuccio, F.P., Surtees, P.G., Luben, R., Brayne, C., Khaw, K.T. (2016). [Daytime Napping, Sleep Duration and Increased 8-year Risk of Type 2 Diabetes in a British Population](#). *Nutrition, Metabolism and Cardiovascular Diseases*, 26(11): 996-1003.
- Lim, M. M., Gerstner, J. R., & Holtzman, D. M. (2014). [The Sleep–Wake Cycle and Alzheimer's Disease: What Do We Know?](#) *Neurodegenerative disease management*, 4(5), 351-362.
- Ohayon, M., Wickwire, E. M., Hirshkowitz, M., Albert, S. M., Avidan, A., Daly, F. J., et al. (2017). [National Sleep Foundation's Sleep Quality Recommendations: First Report](#). *Sleep Health*, 3(1), 6-19.
- Ooms, S., & Ju, Y. E. (2016). [Treatment of Sleep Disorders in Dementia](#). *Current treatment options in neurology*, 18(9), 40.
- Palmer, C. A., & Alfano, C. A. (2017). [Sleep and Emotion Regulation: an Organizing, Integrative Review](#). *Sleep medicine reviews*, 31, 6-16.
- Petrovsky, D. V., McPhillips, M. V., Li, J., Brody, A., Caffee, L., & Hodgson, N. A. (2018). [Sleep Disruption and Quality of Life in Persons with Dementia: A State-of-the-Art Review](#). *Geriatric Nursing*, 39(6), 640-645.
- Smith, T. J., Wilson, M. A., Karl, J. P., Orr, J., Smith, C. D., Cooper, A. D., ... Montain, S. J. (2018). [Impact of Sleep Restriction on Local Immune Response and Skin Barrier Restoration With and Without "Multinutrient" Nutrition Intervention](#). *Journal Of Applied Physiology (Bethesda, Md.: 1985)*, 124(1), 190–200.
- U.S. Department of Health and Human Services, Centers for Medicare & Medicaid Services. (2018). [Frequently Asked Questions Related to Long Term Care Regulations, Survey Process, and Training](#).
- Wennberg, A.M.V., Wu, M.N., Rosenberg, P.B., & Spira, A.P. (2017). [Sleep Disturbance, Cognitive Decline, and Dementia: A Review](#). *Semin Neurol*, 37(4): 395-406.